

CWRU-P22-96

December 1996

Comment on “Constraints on the strength of primordial B-fields from big bang nucleosynthesis reexamined”

Peter J. Kernan, Glenn D. Starkman and Tanmay Vachaspati

Department of Physics
Case Western Reserve University
10900 Euclid Ave., Cleveland, OH 44106-7079

Abstract

Recently Cheng, Olinto, Schramm and Truran [1] reexamined the constraints on the strength of primordial B-fields from big bang nucleosynthesis (BBN). Their bottom line agreed with that of an earlier recent paper on the subject [2], both in its final limit on the B-field during BBN, and in its conclusion that for allowed values of the B-field, the dominant factor for BBN is the increased expansion rate at a given temperature caused by the energy density of the magnetic field, $B^2/8\pi$. However, their conclusion that weak interaction rates increased with increasing B-field at these low field values contradicted the earlier results of [2]. In this comment we point out that the Taylor series expansion of the weak interaction rate about $B = 0$ used in [1] is not well-defined, while the Euler-McLaurin expansion of [2] is well-behaved and reliable. Using the Euler-McLaurin expansion we find that the weak interaction rates decrease rather than increase with increasing B-field at small values of the B-field.

As discussed by Cheng, Olinto, Schramm and Truran [1] “BBN provides a unique quantitative window for processes occurring in the early universe between temperatures of 1 – 0.01MeV. A primeval magnetic field existing during this period would have three major effects on BBN” one of which is that it would alter the weak interaction rates. Equations (2.4-2.6) of [1] give the reaction rates for $ne^+ \rightarrow p\bar{\nu}_e$, $n\nu_e \rightarrow pe^-$ and $n \rightarrow pe^-\nu_e$ respectively. For example the rate for $n\nu \rightarrow pe^-$ is

$$\begin{aligned} \lambda_{n\nu \rightarrow pe^-} &= \frac{g_V^2(1+3\alpha^2)m_e^5\gamma_i T_\nu^2}{4\pi^3 T_i^2} \times \\ &\times [\sum_{n_s=0}^{\infty} [2 - \delta_{n_s 0}] \int_{\sqrt{1+2\gamma n_s + \kappa}}^{\infty} \frac{d\epsilon(\epsilon-\kappa)(\epsilon-q)^2}{\sqrt{(\epsilon-\kappa)^2 - (1+2\gamma n_s)}} \frac{e^{\epsilon Z_e + \phi_e}}{1+e^{\epsilon Z_e + \phi_e}} \frac{1}{1+e^{(\epsilon-q)Z_\nu + \phi_\nu}} \\ &- \sum_{n_s=0}^{n_{smax}+1} [2 - \delta_{n_s 0}] \int_{\sqrt{1+2\gamma n_s + \kappa}}^q \frac{d\epsilon(\epsilon-\kappa)(\epsilon-q)^2}{\sqrt{(\epsilon-\kappa)^2 - (1+2\gamma n_s)}} \frac{e^{\epsilon Z_e + \phi_e}}{1+e^{\epsilon Z_e + \phi_e}} \frac{1}{1+e^{(\epsilon-q)Z_\nu - \phi_\nu}}] \end{aligned} \quad (1)$$

where $\alpha = g_V/g_A, \gamma = B/B_c$ (with $B_c = m_e^2/e$), $\gamma_i = B(T = 1MeV)/B_c, n_s = n + \frac{1}{2} - s_z$ (with n the principal quantum number of the Landau level and $s_z = \pm \frac{1}{2}$ the spins, κ is the anomalous magnetic moment term for an electron in the ground state ($n = 0, s_z = 1/2$), $\epsilon = E_e/m_e, q = (m_n - m_p)/m_e, Z_e = m_e/T_e, Z_\nu = m_\nu/T_\nu, \phi_e = \mu_e/T_e, \phi_\mu = \mu_\mu/T_\mu, m_i$ are the rest masses of species i, T_i are their temperatures and μ_i their chemical potentials. n_{smax} is the largest integer in $[(q - \kappa)^2 - 1]/2\gamma$. In the absence of chemical potentials and ignoring the anomalous electron magnetic moment both of which are lower order effects at the magnetic fields which ultimately prove to be of interest (unless there is a non-standard neutrino chemical potential) this can be rewritten as (equation (22) of [2]):

$$\lambda_{n\nu \rightarrow pe^-} = \frac{G_F^2 T_\gamma^2 (g_V^2 + 3g_A^2)}{(2\pi)^3} z \sum_{n=0}^{\infty} (2 - \delta_{n0}) \int_0^{\infty} dp_z E_\nu^2 g(E_e/T_\gamma) f(E_\nu/T_\nu) \quad (2)$$

where we have rewritten the integral over the electron energy in terms of an integral over the electron momentum parallel to the field, p_z . Here G_F is the Fermi constant, $T_\gamma = T_e$ is the photon temperature, $f(E_\nu/T_\nu)$ is the Fermi-Dirac distribution, and $g = 1 - f$ is the Fermi blocking factor, $z \equiv 2eB/T_\gamma^2$ (so that also $z = 2e\gamma_i \frac{T_\gamma^2}{T_i^2} \frac{m_e^2}{eT_i^2}$). The electron energy and momentum are related by

$$E_e = (p_z^2 + m_e^2 + 2eBn)^{1/2}, \quad (3)$$

while energy conservation gives:

$$E_\nu = E_e - (m_n - m_p) \equiv E_e - \Delta, \quad (4)$$

and $f(E_\nu/T_\nu)$ is taken to be zero if $E_\nu < 0$. We notice that λ is of the form $zh(z)$, where $h(z)$ is a complicated function of z , and so

$$\frac{d\lambda}{dz} = h(z) + z \frac{dh(z)}{dz} \quad (5)$$

We wish to examine the weak field limit $z \rightarrow 0$ in order to understand whether magnetic fields speed-up or slow-down weak interactions. The temptation (see [1]) is to set the second term to zero, and notice that $h(z)$ is the sum of positive terms, and thus conclude that $\frac{d\lambda}{dz} > 0$. The problem is that

$$\lim_{z \rightarrow 0} h(z) = \infty \quad (6)$$

We can see this explicitly by setting $B = 0$ (*i.e.* $\gamma = 0 = \gamma_i$) in (1) and noticing that (with $\kappa = \phi_i = 0$) the argument of the integral is finite, positive and independent of n_s , hence the sum from $n_s = 0, \dots, \infty$ is badly divergent. In fact $h(z) \propto 1/z$ at small z since $\lambda(z) = zh(z)$ should go to the interaction rate in the absence of a magnetic field as we let $z \rightarrow 0$.

The Taylor series of λ about $B = 0$ is therefore ill-defined. However, we can use the Euler-McLaurin expansion for λ ([2] equation (23)):

$$\begin{aligned} \lambda_{n\nu \rightarrow pe^-} &= \frac{G_F^2 T_\gamma^2 (g_V^2 + 3g_A^2)}{(2\pi)^3} \left\{ 2 \int_0^\infty dx \int_{p_{min}(x)}^\infty dp_z G_-(E_e(p_z, x)) \right. \\ &\quad \left. - \frac{z^2 T_\gamma^2}{12} \int_{p_{min}(0)}^\infty dp_z \left[\frac{1}{E_e} \frac{dG_-(E_e)}{dE_e} \right]_{x=0} \right\} + \mathcal{O}(z^3) \end{aligned} \quad (7)$$

where

$$E_e(p_z, x) = (p_z^2 + m_e^2 + x)^{1/2}, \quad (8)$$

$$p_{min}(x) = (\Delta^2 - m_e^2 - xT_\gamma^2)^{1/2} \quad (9)$$

and

$$G_-(E_e) = (E_e - \Delta)^2 \frac{1}{1 + e^{(E_e - \Delta)/T_\nu}} \frac{1}{1 + e^{-E_e/T_\gamma}}. \quad (10)$$

Thus

$$\begin{aligned} \left[\frac{1}{E_e} \frac{dG_-(E_e)}{dE_e} \right] &= \frac{E_e - \Delta}{E_e} \frac{1}{1 + e^{(E_e - \Delta)/T_\nu}} \frac{1}{1 + e^{-E_e/T_\gamma}} \\ &\times \left[2 - \frac{E_e - \Delta}{T_\nu} \frac{1}{1 + e^{-(E_e - \Delta)/T_\nu}} + \frac{E_e}{T_\gamma} \frac{1}{1 + e^{E_e/T_\gamma}} \right] \end{aligned} \quad (11)$$

The terms of the series in (7) are perfectly finite in the limit $z \rightarrow 0$. The zeroth order term recovers exactly the usual $B = 0$ result. Interestingly, we find by explicit calculation that the first order term in B actually vanishes, in apparent contradiction to the positivity of the first order term in the Taylor series. This is of course because the Taylor expansion is not well defined; every term of the Taylor series contributes at arbitrarily small B-field.

The sign of the z^2 term in λ can be seen by inspection of (11). We see that only the second term in the expression in square brackets is negative and that for $x = 0$, by the time E_e is large enough for this term to dominate over the other two terms in the square brackets, the prefactor is exponentially suppressed. The coefficient of the z^2 term of (7) is therefore negative, and the B-field decreases this rate. Similar arguments can be applied to the other $2 \rightarrow 2$ rates. Though the conclusions are not always so clear analytically, numerical studies [2] show a decrease in all the 2-body rates for $B > 0$, in contradiction to the conclusions of [1]. Fortunately as both [1] and [2] agree, this effect is subdominant; the dominant effect of the B-field on nucleosynthesis is simply the B^2 contribution to the energy density, which increases the expansion rate.

References

- [1] B. Cheng, A.V. Olinto, D.N. Schramm and J.W. Truran, Phys. Rev **D54**, 4714 (1996), astro-ph/9606163.
- [2] P. Kernan, G.D. Starkman and T. Vachaspati, Phys. Rev (in press); CWRU-P10-95, astro-ph/9509126 (1995).